

AD-A148 609

NEOHMOGLOBINS AND CROSS-LINKED HEMOGLOBINS AS BLOOD  
SUBSTITUTE(U) MARYLAND UNIV BALTIMORE E BUCCI DEC 82  
DAMD17-81-C-1191

1/1

UNCLASSIFIED


F/G 6/1

NL

END

FILMED

DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A148 609

DTIC FILE COPY

REPORT NUMBER

1

TITLE

Neohemoglobins and Cross-Linked Hemoglobins  
as Blood Substitute

TYPE OF REPORT

ANNUAL SUMMARY REPORT

AUTHOR(s)

ENRICO BUCCI, MD, Ph.D.

Professor of Biochemistry

DECEMBER 1982

Supported by

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND

Fort Detrick, Frederick, Maryland 21701

Contract No. DAMD17-81-C-1191


University of Maryland at Baltimore

Baltimore, Maryland 21201

DOD DISTRIBUTION STATEMENT

Approved for Public Release; Distribution Unlimited

The findings in this report are not to be construed as an official Department of the  
Army position unless so designated by other authorized documents



REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Neohemoglobins and Cross-Linked Hemoglobins as Blood Substitute		5. TYPE OF REPORT & PERIOD COVERED Annual report Sept. 1, 1981 - Nov. 15, 1982
7. AUTHOR(s) Enrico Bucci		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Maryland at Baltimore 660 W. Redwood St., Baltimore, Maryland 21201		8. CONTRACT OR GRANT NUMBER(s) DAMD17-81-C-1191
11. CONTROLLING OFFICE NAME AND ADDRESS Commander, US Army Medical Research and Development Command, ATTN: SGRD-RMS, Fort Detrick, Frederick, MD 21701-5012		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62772A.3S162772A874.AC.139
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE December 1982.
		13. NUMBER OF PAGES 17
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) A: Approved for public release. Distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Hemoglobin Diaspirin Dibromosalicylic-Acid Bromohemoglobin cyano hemoglobin intramolecular cross-link		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We have investigated the effect of heme modifications and specific cross-linking of $\beta$ subunits in human and bovine hemoglobins. It appears that both techniques can be utilized for obtaining products potentially useful as oxygen carriers in resuscitative fluids.		

TITLE

Neohemoglobins and Cross-Linked Hemoglobins  
as Blood Substitute

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A1	

## SUMMARY

Starting from deuteroporphyrin we synthesized 2,4-dibromo,2(or 4)-monocyano and 2-cyano-4-vinyl heme. These hemes were reinserted into apohemoglobin and the oxygen binding properties of the resulting hemoglobins were tested at various pH values between 6 and 8 in 0.05 M Tris or Bis-Tris buffer at 25° C. The oxygen affinity of 2,4-dibromo- and 2-cyano-4-vinyl hemoglobins was very similar to that of normal SFH. The oxygen binding cooperativity gave a value of  $n$  in the Hill plots between 1.8-2.0. The other hemoglobins had a somewhat higher oxygen affinity and a lower cooperativity with values of  $n$  near 1.5.

2,4-Dicyano-heme was also synthesized, however it failed to recombine with apohemoglobin.

Fumaryl, succinyl, and muconyl diaspirins were synthesized and utilized to produce intramolecular cross-linking of the subunits of human and bovine hemoglobins. After 2 hours at 37° C in 0.1 M bis-Tris buffer at pH 7.2, in the presence of 1 mM quantities of the various diaspirins, human hemoglobin reacted very well (70-90%) with the three reagents. Bovine hemoglobin showed a substantial reaction only with fumaryl diaspirin. The oxy, carbonmonoxy and deoxy derivatives of the two hemoglobins were used for the chemical treatment. The extent of cross-linking was tested by SDS gel electrophoresis. This analysis indicated that the major portion of the reacted hemoglobin was cross-linked producing subunits pairs, non dissociable upon denaturation in the presence of detergents.

The oxygen binding properties of these compounds were tested in the same buffers as above described.

For human hemoglobin, cross-linking of the deoxy derivative did not change the oxygen affinity and lowered only a little the binding cooperativity (values of  $n$  near 2.0). Instead, cross-linking of the deoxy derivative produced a compound with an oxygen

affinity lower than that of normal SFH.

For bovine hemoglobin, cross-linking of the oxy and carboxy derivatives increased substantially the oxygen affinity and eliminated the oxygen binding cooperativity. Cross linking of the deoxy derivative left practically unchanged the oxygen affinity and only lowered a little the oxygen binding cooperativity of bovine hemoglobin (values of  $n$  near 2.0).

## TABLE OF CONTENT

	Page
Front Page .....	1
Report Documentation Page .....	2
Title Page .....	3
Summary .....	4
Table of Content .....	6
Report .....	7
References .....	11
Legends .....	12
Figures 1 to 5 .....	13



## REPORT

### MATERIALS AND METHODS

Stroma free-human and bovine hemoglobins were prepared by the filtration method. The respective heme-free proteins (apohemoglobins) were prepared by extraction with methyl-ethylketone (Teale, 1959). Chemical modification of deuteroporphyrins were obtained as described by Caughey et. al. (1966), Falk (1964), and as reported in the monography of Smith (1975). Some derivatives: 2,4-dibromo, 2-formyl-4-vinyl-, and 2 (or 4)-monocyano-deuteroporphyrin-IX-dimethyl ester were obtained from Mid-Century. All of the product prepared and purchased were chromatographically pure. Iron was inserted in the various porphyrins using a large excess of ferrous sulfate on the dimethyl ester derivatives. Recombination of apohemoglobin and neohemes was performed as described by Rossi-Fanelli et. al. (1958).

Acyl derivatives of 3,5-dibromosalicylic acid (diaspirins) were obtained as described by Zaugg et. al., (1980). Cross-linking by diaspirins was performed using a 2 molar excess of the reagent over tetrameric hemoglobin in 0.1 M tris buffer at pH between 7.2 - 7.4, for 2 hours at 37°C. Oxy, carbonmonoxy and deoxy hemoglobins were used. Deoxygenation was obtained adding 1 mg/ml of Na<sup>+</sup> dithionite to solutions previously flushed with nitrogen, in anaerobic environment. The extent of the reaction was followed by electrophoresis with the Beckman microzone system. Acylated hemoglobins had a faster anodic mobility. Polyacrylamide SDS gel electrophoresis (Walder et al, 1979) was used for monitoring the amount of cross-linked protein.

Recombined neohemoglobins and cross-linked hemoglobins were purified by chromatography on CM cellulose using a linear gradient formed by equal volumes of 0.01 M phosphate buffer at pH 6.2 and 0.04 M dibasic phosphate.

Densitometric scans of polyacrylamide gels and of microzone strips were performed with a Joyce and Loeb microdensitometer. Sedimentation velocities were measured in a Beckman model E analytical ultracentrifuge in 0.15 M tris buffer at pH 7.4, at temperatures between 18 and 20°C. Oxygen equilibria were measured using a Gill's cell apparatus (Dolman and Gill, 1978) so to scan the absorption spectrum of the solutions at each step of oxygenation. Ferric hemoglobin formation was less than 5% of the total protein. The hemoglobin solutions were at concentrations between 5 and 10%. Temperature was controlled with a Digitek 500 digital thermometer.

## RESULTS

### Neohemoglobins

Fig. 1 shows the structure of heme where positions 2 and 4 are occupied by vinyl residues. These positions were variously substituted with electron withdrawing groups so to reduce the affinity of the iron in the heme for oxygen.

Neohemes used were 2,4-dibromo-, 2 (or 4)-mono-cyano-(the mixture of the 2 and 4 substituted hemes had not been resolved), and 2-cyano-4-vinyl-heme. In the mono-cyano-derivative the respective 4 and 2 positions were occupied by a proton.

Figs. 2 and 3 show the oxygen affinity of the neohemoglobins as compared to that on normal human SFH either in 0.05 M phosphate, or in 0.05 M Tris or in 0.05 M Bis-tris buffers at various pH values, at 25°C, in the absence of CO<sub>2</sub>. It appears that in human hemoglobin the various neohemes tested were able at best to reproduce the oxygen affinity of untreated hemoglobin, with a lower cooperativity.

Figs. 2 and 3 also shows that there was specie specificity in the interaction of apohemoglobins with neohemes. In fact 2,4-dibromo-heme drastically increased the oxygen affinity of bovine hemoglobin, which normally is even lower than that of human SHF. Also it abolishes the heme-heme interaction. In horse hemoglobin 2,4-dibromo-heme appears to produce a hemoglobin with a lower oxygen affinity than human SHF. Untreated horse and human hemoglobins have identical oxygen affinities.

### Cross-Linked Human and Bovine Hemoglobins

For this reaction we have used the bio-3,5-dibromo-salicylic derivatives of fumaryl, succinyl and muconyl residues, whose structural formulas are illustrated in fig 4.

The extent of reaction, estimated from microzone electrophoresis, was very high for human hemoglobin, reaching the 80-90% of the total for both the liganded and unliganded derivatives. Bovine hemoglobin was less reactive. The fumaryl-diaspirin produced a reaction of 60-70%. The succinyl and muconyl diaspirins produced only 30% reaction or less.

The extent of cross-linking estimated by SDS gel electrophoresis indicated that in non-purified samples the extent of cross-linking was less, about 2/3, than that expected from the extent of the reaction with the diaspirins. After purification by chromatography this discrepancy was corrected and the amount of cross-linked protein approached the 50% mark expected from cross-linking of the subunits only as described by Walder et. al. (1980).

Native and cross-linked samples of human and bovine hemoglobins were indistinguishable in sedimentation velocity experiments which in any case produced single symmetrical peaks with a sedimentation constant near  $S_{20,w} = 4.2$ , independent of protein concentration.

Functional tests of cross-linked hemoglobins are presented in Fig. 5. Circles refer to human and triangles to bovine hemoglobin. Closed symbols imply cross-linking of the oxy-derivatives. For human hemoglobin, hemoglobin cross-linking of the oxy derivative left the oxygen affinity practically unchanged, while cross-linking of the deoxy

derivative produced a substantial decrease of the oxygen affinity. The value of  $n$  in the Hill plots was near 2.7 for native hemoglobin and between 1.6 and 1.9 for cross-linked non purified samples. For bovine hemo globin cross-linking of the oxy derivative produced a substantial increase of the oxygen affinity, while in the deoxy derivative it left the oxygen affinity practically unchanged. The values of  $n$  in the Hill plots were near 2.8 for the native protein and between 1.5 and 1.9 for unpurified samples.

Purification by chromatography of human or bovine hemoglobins cross-linked as deoxy derivative, seemed to increase the value of  $n$  in the functional tests.

## CONCLUSIONS

The presence of the porphyrin ring of the heme of electron withdrawing groups is expected to lower the affinity of the iron atom for ligands. The neohemes here investigated show that at least in human hemoglobin the mere substitution of the vinyl groups in position 2 and 4 with other electron withdrawing agents is not enough for reducing the oxygen affinity of the protein to levels desirable in blood substitutes.

In regard to cross-linking with diasprinis, the decreased oxygen affinity of human hemoglobin, when reacted as unliganded derivative, is extremely interesting and promising.

The drastic increase of the oxygen affinity of bovine hemoglobin upon cross-linking of its liganded form, implies a distortion of the molecule, which is not produced when the reaction is performed on the deoxy derivative. In this case the oxygen binding cooperativity is decreased, however it is still present and the oxygen affinity remains lower than that of human SHF, making the protein suitable as blood substitute. The tetrameric structure of the proteins stabilized by cross-linking should allow a longer retention time after transfusion than for untreated human SHF.


## REFERENCES


- Caughey WS, Fujimoto WJ, Johnson B (1966). Substituted Deuteroporphyrins. *Biochemistry* 5:3830
- Dolman D, Gill SJ (1978). Membrane Covered Thin-Layer Optical Cell for Gas Reaction Studies of Hemoglobin. *Anal Biochem* 87:127
- Falk, JE (1964). "Porphyrins and Metalloporphyrins". New York, Amsterdam: Elsevier Pub Co
- Rossi Fanelli A, Antonini A, Caputo A (1958). Studies on the Structure of Hemoglobin. I Physicochemical Properties of Human Globin. *Biochim Biophys Acta* 30:608
- Smith KM (1975). "Porphyrins and Metalloporphyrins". New York, Amsterdam: Elsevier Pub Co
- Teale FWJ (1959). Cleavage of the Haem-Protein Link by Acid Methylethylketone. *Biochim Biophys Acta* 35:543
- Walder JA, Zaugg RH, Walder RY, Steele JM, Klotz IM (1979). Diaspirins That Cross-Link Chains of Hemoglobin. *Biochemistry* 18:4265
- Walder JA, Walder RY, Arnone A (1980). Development of Antisickling Compounds That Chemically Modify Hemoglobin S Specifically within the 2,3-Diphosphoglycerate Binding Site. *J Mol Biol* 141:195.

## LEGENDS

- Fig. 1      Ferroprotoporphyrin-IX
- Fig. 2      Oxygen binding characteristics of the indicated reconstituted hemoglobin in 0.05 M Tris or Bis-Tris buffers at 25°C in the presence of 5% CO<sub>2</sub>. The symbol A refers to human SFH.
- Fig. 3      Oxygen binding characteristics of the indicated reconstituted hemoglobins in 0.05 M phosphate buffers at 25°C, in the presence of 5% CO<sub>2</sub>. The symbol A refers to human SFH.
- Fig. 4      Various esters of bis-3,5-dibromosalicylic acid.
- Fig. 5      Oxygen binding characteristics of various cross-linked hemoglobins. A refers to normal human SFH and B to normal bovine SFH. Δ, ▲)Bovine hemoglobin, O, ●)human hemoglobins. Open symbols refer to cross-linking reactions performed on the deoxyderivatives of human or bovine hemoglobin.

Fig.1













Fig. 4



BIS-3,5-DIBROMOSALICYLIC ACID





**END**

**FILMED**

**2-85**

**DTIC**